

# Using $K_\alpha$ images to characterize pre-plasma and electron refluxing in intense laser experiments


**V.M. Ovchinnikov**  
**The Ohio State University**

2011 Fall Meeting of the DPP APS  
Salt Lake City, UT  
November 17, 2011

# Collaborators



K.U. Akli, R.R. Freeman, G.E. Kemp, A. Krygier,  
D.W. Schumacher, L. Van Woerkom



R. Stephens



F.N. Beg, S. Chawla, D.P. Higginson, C. Jarrot, C. Murphy,  
H. Sawada, M.S. Wei, B. Westover, T. Yabuuchi



C.D. Chen, D.S. Hey, M.H. Key, A.J. Mackinnon,  
H.S. MacLean, A.G. MacPhee, A. Link, P.K. Patel



R. Fedosejevs, H. Friesen, Y. Tsui



# Motivation


- Angular distribution of laser generated hot electrons is crucial.
- Recent experimental measures of the divergence vary widely.
- $K_\alpha$  imaging is a primary experimental measure.
- PIC simulations tend to indicate a large *intrinsic* divergence in the LPI region ( $\Theta_{1/2} = 50-60^\circ$ ).



**This talk will show how a large intrinsic electron divergence can result in varying divergences as measured by  $K_\alpha$  imaging.**

# We've shot slabs and buried cone targets

## Cone-guided Fast Ignition:




*Buried cone target with a Get-Lost Layer (GLL)*

*$K_\alpha$  image of a buried cone target*


# Recent Titan Results: $K_{\alpha}$ imaging

## 2009 Run: Buried Cone Results. $\left[ \frac{W}{\lambda} \right]_{1/2} = 40-45^{\circ}$



Detector used:  
X-ray CCD

## 2010 Run: Buried Cones Results. $\left[ \frac{W}{\lambda} \right]_{1/2} = 12-15^{\circ}$



Detector used:  
Image Plates

# Simulation setup for the 2009 experiment

Fully kinetic simulations, using modified PIC code LSP

## (1) TARGET

Buried cone target  
(30° cone angle)  
Exponential preplasma

## (2) GRID


2D Cartesian (XZ)  
Cell size: from  $\lambda/8$  to  $\lambda$

## (3) LASER

$\lambda = 1 \mu\text{m}$   
 $I = 1 \times 10^{20} \text{ W/cm}^2$   
 $w_0 = 7 \mu\text{m}$   
sine<sup>2</sup> envelope w/  
 $\tau = 700 \text{ fs}$   
propagates along +x  
z-polarized

## (4) RUN DURATION

$\sim 20 \text{ ps}$   
with  $\Delta t = 30$  timesteps/optical cycle




Preplasma scale length is the only free parameter


# Simulation vs. 2009 Experiment: $L = 5 \mu\text{m}$

Red – simulation  
Grey - experiment

a) fluor depth:  $10 \mu\text{m}$




b) fluor depth:  $100 \mu\text{m}$



Preplasma scale length  $L = 5 \mu\text{m}$

# Simulation vs. 2009 Experiment: $L = 2.5 \mu\text{m}$


Red – simulation  
Grey - experiment



Preplasma scale length  $L = 2.5 \mu\text{m}$

# Simulation vs. 2009 Experiment: $L = 3.75 \mu\text{m}$

Red – simulation  
Grey - experiment




Preplasma scale length  $L = 3.75 \mu\text{m}$


# Effect of the Get Lost Layer

Red – simulation  
Grey - experiment

a) fluor depth: 10  $\mu\text{m}$





b) fluor depth: 100  $\mu\text{m}$




Preplasma scale length  $L = 3.75$   $\mu\text{m}$  (no GLL)

# Simulation vs. experiment for a slab with no GLL




Slab target, no GLL


# Experimental results for 2010 experiment






$$\varpi_{1/2} = 12-15^\circ$$






# Noise issue in IP-based images






# Simulations for 2010 experiment




Red – simulation  
Grey - experiment



Preplasma  
scale  
length:  
 $L=2.5 \mu\text{m}$





# Slab targets with GLL




# Slab results

15  $\mu\text{m}$




Red – simulation  
Grey - experiment

100  $\mu\text{m}$



Preplasma  
scale  
length:  
 $L = 1.75 \mu\text{m}$

200  $\mu\text{m}$



# Summary



- Good match between experiment and simulation.
- Wide range of targets using data collected with 2 different detector types over 2 years.

We find that  $K_\alpha$  images

- provide a poor measure of intrinsic divergence.
- yield a bulk divergence angle that is low compared to intrinsic.
- can characterize the pre-plasma near the critical surface.
- can be used to check the performance of GLL's.

**$K_\alpha$  images can still be used to benchmark PIC codes from which the intrinsic divergence can be determined.**

Thank you!